



# Allometric Equations for Perennial Grasses in the Desert Grassland

Aleta Nafus, Mitchel P. McClaran, Chad McMurtry, School of Natural Resources, University of Arizona, Tucson Arizona 85721

## Introduction

- Productivity and amount of standing biomass are critical to understand ecosystem function and response to management.
- Non-destructive estimates of biomass reduce the time investment and impact of destructive estimates.
- Allometric relationships provide non-destructive estimates of biomass based on size and mass relationships.
- Diameter is most strongly correlated with biomass although height and grazing history should improve the model.
- Allometric relationships should differ between plants with/without long exposure to livestock grazing because repeated defoliation 1) removes standing biomass and 2) creates prostrate growth through increased tillering
- Therefore, we expect exposure to grazing to 1) reduce the total standing mass-size<sup>-1</sup> relationship and 2) increase the productivity mass-size<sup>-1</sup> relationship if increased tillering occurs.

## Objectives

- Generate the allometric relationships between total mass and productivity (green mass) for 8 common grass species in the Desert Grassland.
- Compare the allometric relationships between plants exposed to different livestock grazing histories.

## Methods

- Santa Rita Experimental Range (Pasture 2SW) grazed moderately since 1950, but ungrazed during current summer grazing season, and adjacent Rodent Station Exclosure ungrazed since 1903.
- 25-45 plants species<sup>-1</sup> along a size (diameter) gradient collected in August 2005
- 8 species (species groups)
- Aristida* species (ARI), *Bouteloua eriopoda* (BOER), *Digitaria californica* (DICA); *Eragrostis curvula* (ERCU), *E.lehmanniana* (ERLE), *Heteropogon contortus* (HECO), *Muhlenbergia porteri* (MUPO) and *Setaria leucopila* (SELE).
- HECO and BOER collected in ungrazed area only.

## Measurements

- Plant Mass:  $Y_{Total}$  = total mass (g plant<sup>-1</sup>) and  $Y_{Green}$  = total green mass (g plant<sup>-1</sup>).
- Plant Size: D=basal diameter (cm) and H=tallest leaf collar (cm) (Figure 2).
- Grazing History (G): where 0 = ungrazed and 1=grazed, expressed as interaction with size (D\*G and H\*G).

## Model Development

- Natural log transformation of mass and size to normalize residuals.
- Forward Stepwise Regression with  $p < 0.05$  for entry.

## Results




Figure 1. *Eragrostis curvula* showing effect of grazing on regression models for grazed and ungrazed mass by basal diameter

Table 1: Peak Seasonal Production (Summer Production) ( $Y_{Green}$ )

Species	Model Parameters							R2	SEE
	Allometric models	Intercept	Diameter	Height	Diam* Grazing	Height* Grazing	n		
ARI	D+(D*G)	1.032	1.757		-0.687		28	0.878	0.476
BOER	D	0.448	1.335				22	0.855	0.521
DICA	D+H	-3.478	0.92	1.323			36	0.895	0.359
ERCU	D+(D*G)	0.885	1.136		-0.379		29	0.865	0.483
ERLE	D+H+(D*G)	-4.877	0.938	1.696	-0.225		35	0.960	0.382
HECO	D+H	-0.812	0.915	0.716			20	0.931	0.384
MUPO	D+H	-5.81	0.863	2.214			34	0.917	0.453
SELE	D+H+(H*G)	-1.601	1.22	0.809		-0.084	36	0.928	0.419

Table 2: Total Standing Biomass ( $Y_{Total}$ )

Species	Model Parameters							R2	SEE
	Allometric models	Intercept	Diameter	Height	Diam* Grazing	Height* Grazing	n		
ARI	D+D*G	1.101	1.798		-0.683		28	0.877	0.472
BOER	D	0.813	1.473				22	0.900	0.466
DICA	D+H+(D*G)	-3.816	1.272	1.416	-0.244		36	0.946	0.323
ERCU	D+(D*G)	1.093	1.282		-0.407		29	0.913	0.427
ERLE	D+H	-4.877	0.838	1.726			35	0.954	0.404
HECO	D+H	-0.812	0.915	0.716			20	0.937	0.388
MUPO	D+H	-4.93	0.97	2.046			34	0.912	0.494
SELE	D+H+(H*G)	-1.366	1.269	0.771		-0.112	36	0.934	0.411

## Results and Discussion

### Objective 1

- Allometric models had strong  $R^2$  values ( $> 0.88$  and  $0.86$ ) and high confidence levels ( $SEE < 0.49$  and  $0.52$ ) for Total Mass ( $Y_{Total}$ ) and Summer Productivity ( $Y_{Green}$ ), respectively (Table 1 and Table 2).
- Diameter (D) was the most significant explanatory variable, and height was included as an explanatory variable included in most models (Table 1 and Table 2).

### Objective 2:

- As expected, grazing history reduced the mass
- size<sup>-1</sup> relationship for Total Mass ( $Y_{Total}$ ), but this occurred in only 4 of 6 species (ARIS, DICA, ERCU, and SEMA; Table 2).
- Grazing history reduced the mass size<sup>-1</sup> relationship for Green Mass ( $Y_{Green}$ ) in 4 of 6 species (ARIS, ERCU, ERLE, and SEMA; Table 1, Figure 1).
- Reduction in the Green Mass size<sup>-1</sup> relationship suggests that a long exposure to defoliation resulted in a concentration of mass in prostrate growth. This is especially apparent in SEMA where the grazing interaction was with height.

## Implications

- Models can be used to accurately predict both peak seasonal biomass and total standing biomass on a site based on morphological measures of basal diameter and height and non-morphological site factors like grazing history.
- Grazing history is a significant explanatory variable for at least some species when creating allometric models and should be considered when attempting to predict biomass on a site

### Acknowledgements

Partial funding provided by USDA-NRI Ecosystem Management Program and Arizona Agricultural Experiment Station. Thanks to Dr. Steven Archer and Dr. Heather Throop for obtaining the funding. Lab help was provided by M. Castillo and K. Lee. Field help was provided by S. Archer, E. Hurd, F. Mashiri, and J. Mudd.



Figure 2: Basal diameter and height measurement